

Errors for Linear Systems

When we solve a linear system $Ax = b$ we often do not know A and b exactly, but have only approximations \hat{A} and \hat{b} available. Then the best thing we can do is to solve $\hat{A}\hat{x} = \hat{b}$ exactly which gives a different solution vector \hat{x} . We would like to know how the errors of \hat{A} and \hat{b} influence the error in \hat{x} .

Example: Consider the linear system $Ax = b$ with

$$\begin{bmatrix} 1.01 & .99 \\ .99 & 1.01 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}.$$

We can easily see that the solution is $x = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$. Now let us use the slightly different right hand side vector $\hat{b} = \begin{bmatrix} 2.02 \\ 1.98 \end{bmatrix}$ and solve the linear system $A\hat{x} = \hat{b}$. This gives the solution vector $\hat{x} = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$. In this case a small change in the right hand side vector has caused a large change in the solution vector.

Vector norms

In order to measure errors in vectors by a single number we use a so-called *vector norm*.

A **vector norm** $\|x\|$ measures the size of a vector $x \in \mathbb{R}^n$ by a nonnegative number and has the following properties

$$\|x\| = 0 \quad \Rightarrow \quad x = 0$$

$$\|\alpha x\| = |\alpha| \|x\|$$

$$\|x + y\| \leq \|x\| + \|y\|$$

for any $x, y \in \mathbb{R}^n$, $\alpha \in \mathbb{R}$. There are many possible vector norms. We will use the three norms $\|x\|_1$, $\|x\|_2$, $\|x\|_\infty$ defined by

$$\|x\|_1 = |x_1| + \dots + |x_n|$$

$$\|x\|_2 = \left(|x_1|^2 + \dots + |x_n|^2 \right)^{1/2}$$

$$\|x\|_\infty = \max\{|x_1|, \dots, |x_d|\}$$

If we write $\|x\|$ in an equation without any subscript, then the equation is valid for all three norms (using the same norm everywhere).

If the exact vector is x and the approximation is \hat{x} we can define the **relative error with respect to a vector norm** as $\frac{\|\hat{x} - x\|}{\|x\|}$.

Example: Note that in the above example we have $\frac{\|\hat{b} - b\|_\infty}{\|b\|_\infty} = 0.01$, but $\frac{\|\hat{x} - x\|_\infty}{\|x\|_\infty} = 1$. That means that the relative error of the solution is 100 times as large as the relative error in the given data, i.e., the condition number of the problem is at least 100.

Matrix norms

A *matrix norm* $\|A\|$ measures the size of a matrix $A \in \mathbb{R}^{n \times n}$ by a nonnegative number. We would like to have the property

$$\|Ax\| \leq \|A\| \|x\| \quad \text{for all } x \in \mathbb{R}^n \quad (1)$$

where $\|x\|$ is one of the above vector norms $\|x\|_1$, $\|x\|_2$, $\|x\|_\infty$. We define $\|A\|$ as the smallest number satisfying (1):

$$\|A\| := \sup_{\substack{x \in \mathbb{R}^n \\ x \neq 0}} \frac{\|Ax\|}{\|x\|} = \max_{\substack{x \in \mathbb{R}^n \\ \|x\|=1}} \|Ax\|$$

By using the 1, 2, ∞ vector norm in this definition we obtain the matrix norms $\|A\|_1$, $\|A\|_2$, $\|A\|_\infty$ (which are in general different numbers). It turns out that $\|A\|_1$ and $\|A\|_\infty$ are easy to compute:

Theorem:

$$\|A\|_\infty = \max_{i=1,\dots,n} \sum_{j=1,\dots,n} |a_{ij}| \quad (\text{maximum of row sums of absolute values})$$

$$\|A\|_1 = \max_{j=1,\dots,n} \sum_{i=1,\dots,n} |a_{ij}| \quad (\text{maximum of column sums of absolute values})$$

Proof: For the infinity norm we have

$$\|Ax\|_\infty = \max_i \left| \sum_j a_{ij} x_j \right| \leq \max_i \sum_j |a_{ij}| |x_j| \leq \left(\max_i \sum_j |a_{ij}| \right) \|x\|_\infty$$

implying $\|A\|_\infty \leq \max_i \sum_j |a_{ij}|$. Let i_* be the index where the maximum occurs and define $x_j = \text{sign } a_{i_* j}$, then $\|x\|_\infty = 1$ and $\|Ax\|_\infty = \max_i \sum_j |a_{ij}|$.

For the 1-norm we have

$$\|Ax\|_1 = \sum_i \left| \sum_j a_{ij} x_j \right| \leq \sum_j \left(\sum_i |a_{ij}| \right) |x_j| \leq \left(\max_j \sum_i |a_{ij}| \right) \|x\|_1$$

implying $\|A\|_1 \leq \max_j \sum_i |a_{ij}|$. Let j_* be the index where the maximum occurs and define $x_{j_*} = 1$ and $x_j = 0$ for $j \neq j_*$, then $\|x\|_1 = 1$ and $\|Ax\|_1 = \max_j \sum_i |a_{ij}|$. \square

We will not use $\|A\|_2$ since it is more complicated to compute (it involves eigenvalues).

Note that for $A, B \in \mathbb{R}^{n \times n}$ we have $\|AB\| \leq \|A\| \|B\|$ since

$$\|ABx\| \leq \|A\| \|Bx\| = \|A\| \|B\| \|x\|.$$

The following results about matrix norms will be useful later:

Lemma 1: Let $A \in \mathbb{R}^{n \times n}$ be nonsingular and $E \in \mathbb{R}^{n \times n}$. Then $\|E\| < \frac{1}{\|A^{-1}\|}$ implies that $A + E$ is nonsingular.

Proof: Assume that $A + E$ is singular. Then there exists a nonzero $x \in \mathbb{R}^n$ such that $(A + E)x = 0$ and hence

$$\frac{\|x\|}{\|A^{-1}\|} \leq \|Ax\| = \|Ex\| \leq \|E\| \|x\|$$

The left inequality for $b := Ax$ follows from $\|x\| = \|A^{-1}b\| \leq \|A^{-1}\| \|b\|$. As $\|x\| > 0$ we obtain $\frac{1}{\|A^{-1}\|} \leq \|E\|$. \square

Lemma 2: For given vectors $x, y \in \mathbb{R}^n$ with $x \neq 0$ there exists a matrix $E \in \mathbb{R}^{n \times n}$ with $Ex = y$ and $\|E\| = \frac{\|y\|}{\|x\|}$.

Proof: For the infinity-norm we have $\|x\|_\infty = |x_j|$ for some j . Let $a \in \mathbb{R}^n$ be the vector with $a_j = 1$, $a_k = 0$ for $j \neq k$ and let

$$E = \frac{1}{\|x\|} y a^\top,$$

then (i) $a^\top x = \|x\|_\infty$ implies $Ex = y$ and (ii) $\|y a^\top v\|_\infty = |a^\top v| \|y\|_\infty$ with $|a^\top v| \leq \|v\|_\infty$ implies $\|E\| \leq \frac{\|y\|}{\|x\|}$.

For the 1-norm we use $a \in \mathbb{R}^n$ with $a_j = \text{sign}(x_j)$ since $a^\top x = \|x\|_1$ and $|a^\top v| \leq \|v\|_1$.

For the 2-norm we use $a = x / \|x\|_2$ since $a^\top x = \|x\|_2$ and $|a^\top v| \leq \|a\|_2 \|v\|_2 = \|v\|_2$. \square

Condition numbers

Let x denote the solution vector of the linear system $Ax = b$. If we choose a slightly different right hand side vector \hat{b} then we obtain a different solution vector \hat{x} satisfying $A\hat{x} = \hat{b}$. We want to know how the relative error $\|\hat{b} - b\| / \|b\|$ influences the relative error $\|\hat{x} - x\| / \|x\|$ ("error propagation"). We have $A(\hat{x} - x) = \hat{b} - b$ and therefore

$$\|\hat{x} - x\| = \|A^{-1}(\hat{b} - b)\| \leq \|A^{-1}\| \|\hat{b} - b\|.$$

On the other hand we have $\|b\| = \|Ax\| \leq \|A\| \|x\|$. Combining this we obtain

$$\frac{\|\hat{x} - x\|}{\|x\|} \leq \|A\| \|A^{-1}\| \frac{\|\hat{b} - b\|}{\|b\|}.$$

The number $\text{cond}(A) := \|A\| \|A^{-1}\|$ is called **condition number** of the matrix A . It determines how much the relative error of the right hand side vector can be amplified. The condition number depends on the choice of the matrix norm: In general $\text{cond}_1(A) := \|A\|_1 \|A^{-1}\|_1$ and $\text{cond}_\infty(A) := \|A\|_\infty \|A^{-1}\|_\infty$ are different numbers.

Example: In the above example we have

$$\text{cond}_\infty(A) = \|A\|_\infty \|A^{-1}\|_\infty = \left\| \begin{bmatrix} 1.01 & .99 \\ .99 & 1.01 \end{bmatrix} \right\|_\infty \left\| \begin{bmatrix} 25.25 & -24.75 \\ -24.75 & 25.25 \end{bmatrix} \right\|_\infty = 2 \cdot 50 = 100$$

and therefore

$$\frac{\|\hat{x} - x\|}{\|x\|} \leq 100 \frac{\|\hat{b} - b\|}{\|b\|}$$

which is consistent with our results above (b and \hat{b} were chosen so that the worst possible error magnification occurs).

The fact that the matrix A in our example has a large condition number is related to the fact that A is close to the singular matrix $B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$.

The following result shows that $\frac{1}{\text{cond}(A)}$ indicates how close A is to a singular matrix:

Theorem: $\min_{B \in \mathbb{R}^{n \times n}, B \text{ singular}} \frac{\|A - B\|}{\|A\|} = \frac{1}{\text{cond}(A)}$

Proof: (1) Lemma 1 shows: B singular implies $\|A - B\| \geq \frac{1}{\|A^{-1}\|}$.

(2) By the definition of $\|A^{-1}\|$ there exist $x, y \in \mathbb{R}^n$ such that $x = A^{-1}y$ and $\|A^{-1}\| = \frac{\|x\|}{\|y\|}$. By Lemma 2 there exists a matrix $E \in \mathbb{R}^{n \times n}$ such that $Ex = y$ and $\|E\| = \frac{\|y\|}{\|x\|}$. Then $B := A - E$ satisfies $Bx = Ax - Ex = y - y = 0$, hence B is singular and $\|A - B\| = \|E\| = \frac{1}{\|A^{-1}\|}$. \square

Example: The matrix $A = \begin{bmatrix} 1.01 & .99 \\ .99 & 1.01 \end{bmatrix}$ is close to the singular matrix $B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ so that $\frac{\|A - B\|_\infty}{\|A\|_\infty} = \frac{.02}{2} = .01$. By the theorem we have that $0.01 \leq \frac{1}{\text{cond}_\infty(A)}$ or $\text{cond}_\infty(A) \geq 100$. As we say above we have $\text{cond}_\infty(A) = 100$, i.e., the matrix B is really the closest singular matrix to the matrix A .

When we solve a linear system $Ax = b$ we have to store the entries of A and b in the computer, yielding a matrix \hat{A} with rounded entries $\hat{a}_{ij} = fl(a_{ij})$ and a rounded right hand side vector \hat{b} . If the original matrix A is singular then the linear system has no solution or infinitely many solutions, so that any computed solution is meaningless. How can we recognize this on a computer? Note that the matrix \hat{A} which the computer uses may no longer be singular.

Answer: We should compute (or at least estimate) $\text{cond}(\hat{A})$. If $\text{cond}(\hat{A}) < \frac{1}{\varepsilon_M}$ then we can guarantee that *any* matrix A which is rounded to \hat{A} must be nonsingular: $|\hat{a}_{ij} - a_{ij}| \leq \varepsilon_M |a_{ij}|$ implies $\|\hat{A} - A\| \leq \varepsilon_M \|A\|$ for the infinity or 1-norm. Therefore $\frac{\|\hat{A} - A\|}{\|\hat{A}\|} \leq \frac{\varepsilon_M}{1 - \varepsilon_M} \approx \varepsilon_M$ and $\text{cond}(\hat{A}) < \frac{1 - \varepsilon_M}{\varepsilon_M} \approx \frac{1}{\varepsilon_M}$ imply $\frac{\|\hat{A} - A\|}{\|\hat{A}\|} < \frac{1}{\text{cond}(\hat{A})}$. Hence the matrix A must be nonsingular by the theorem.

Now we assume that we perturb both the right hand side vector b and the matrix A :

Theorem: Assume $Ax = b$ and $\hat{A}\hat{x} = \hat{b}$. If A is nonsingular and $\|\hat{A} - A\| < 1/\|A^{-1}\|$ there holds

$$\frac{\|\hat{x} - x\|}{\|x\|} \leq \frac{\text{cond}(A)}{1 - \text{cond}(A) \frac{\|\hat{A} - A\|}{\|A\|}} \left(\frac{\|\hat{b} - b\|}{\|b\|} + \frac{\|\hat{A} - A\|}{\|A\|} \right)$$

Proof: Let $E = \hat{A} - A$, hence $A\hat{x} = \hat{b} - E\hat{x}$. Subtracting $Ax = b$ gives $A(\hat{x} - x) = (\hat{b} - b) - E\hat{x}$ and therefore

$$\begin{aligned}\|\hat{x} - x\| &\leq \|A^{-1}\| \left(\|\hat{b} - b\| + \|E\| \|\hat{x}\| \right) \\ &\leq \|x\| + \|\hat{x} - x\| \\ (1 - \|A^{-1}\| \|E\|) \|\hat{x} - x\| &\leq \|A^{-1}\| \left(\|\hat{b} - b\| + \|E\| \|x\| \right)\end{aligned}$$

Dividing by $\|x\|$ and using $\|b\| \leq \|A\| \|x\| \iff \|x\| \geq \|b\| / \|A\|$ gives

$$\begin{aligned}\left(1 - \|A^{-1}\| \|E\|\right) \frac{\|\hat{x} - x\|}{\|x\|} &\leq \|A^{-1}\| \left(\frac{\|\hat{b} - b\|}{\|b\| / \|A\|} + \|E\| \right) \\ \left(1 - \text{cond}(A) \frac{\|E\|}{\|A\|}\right) \frac{\|\hat{x} - x\|}{\|x\|} &\leq \text{cond}(A) \left(\frac{\|\hat{b} - b\|}{\|b\|} + \frac{\|E\|}{\|A\|} \right)\end{aligned}$$

If $\|E\| < 1 / \|A^{-1}\| \iff \frac{\|E\|}{\|A\|} < 1 / \text{cond}(A)$ we can divide by $\left(1 - \text{cond}(A) \frac{\|E\|}{\|A\|}\right)$. \square

If $\text{cond}(A) \frac{\|\hat{A} - A\|}{\|A\|} \ll 1$ we have that both the relative error in the right hand side vector and in the matrix are magnified by $\text{cond}(A)$.

If $\text{cond}(A) \frac{\|\hat{A} - A\|}{\|A\|} = \|A^{-1}\| \|\hat{A} - A\| \geq 1$ then by the theorem for the condition number the matrix \hat{A} may actually be singular, so that the solution \hat{x} is no longer well defined.

Computing the condition number

We have seen that the condition number is very useful: It tells us what accuracy we can expect for the solution, and how close our matrix is to a singular matrix.

In order to compute the condition number we have to find A^{-1} . This takes $n^3 + O(n^2)$ operations, compared with $\frac{n^3}{3} + O(n^2)$ operations for the LU-decomposition. Therefore the computation of the condition number would make the solution of a linear system 3 times as expensive. For large problems this is not reasonable.

However, we do not need to compute the condition number with full machine accuracy. Just knowing the order of magnitude is sufficient. Since $\|A^{-1}\| = \max_{\|z\|=1} \|A^{-1}z\|$ we can pick any vector z with $\|z\| = 1$ and have the lower bound $\|A\| \geq \|A^{-1}z\|$. Note that $u = A^{-1}z$ is the solution of the linear system $Au = z$. If we already have L, U, p we can easily find u using $u = U \setminus (L \setminus b(p))$ with a cost of only $n^2 + O(n)$ operations. The trick is to pick z with $\|z\| = 1$ such that $\|A^{-1}z\|$ becomes as large as possible, so that it is close to $\|A^{-1}\|$. There are heuristic methods available which achieve fairly good lower bounds: **(i)** Pick $c = (\pm 1, \dots, \pm 1)$ and pick the signs so that the forward substitution gives a large vector, **(ii)** picking $\tilde{c} := z$ and solve $A\tilde{z} = \tilde{c}$ often improves the lower bound. The Matlab functions `cond` and `1/rcond` use similar ideas to give lower bounds for $\text{cond}_1(A)$. Typically they give an estimated condition number c with $c \leq \text{cond}_1(A) \leq 3c$ and require the solution of 2 or 3 linear systems which costs $O(n^2)$ operations if the LU decomposition is known. (However, the Matlab commands `cond` and `rcond` only use the matrix A as an input value, so they have to compute the LU decomposition of A first and need $\frac{n^3}{3} + O(n^2)$ operations.)

Computation in machine arithmetic and residuals

When we run Gaussian elimination on a computer each single operation causes some roundoff error, and instead of the exact solution x of a linear system we only get an approximation \hat{x} . As explained above we should select the pivot candidate with the largest absolute value to avoid unnecessary subtractive cancellation, and this usually is a numerically stable algorithm. However, there is no theorem which guarantees this for partial pivoting (row interchanges). (For “full pivoting” with row and column interchanges some theoretical results exist. However, this algorithm is more expensive, and for all practical examples partial pivoting seems to work fine.)

Question 1: How much error do we have to accept for $\frac{\|\hat{x}-x\|}{\|x\|}$? This is the *unavoidable error* which occurs even for an ideal algorithm where we only round the input values and the output value to machine accuracy, and use infinite accuracy for all computations.

When we want to solve $Ax = b$ we have to store the entries of A, b in the computer, yielding a matrix \hat{A} and a right hand side vector \hat{b} of machine numbers so that $\frac{\|\hat{A}-A\|}{\|A\|} \leq \varepsilon_M$ and $\frac{\|\hat{b}-b\|}{\|b\|} \leq \varepsilon_M$. An ideal algorithm would then try to solve this linear system exactly, i.e., compute a vector \hat{x} such that $\hat{A}\hat{x} = \hat{b}$. Then we have

$$\frac{\|\hat{x}-x\|}{\|x\|} \leq \frac{\text{cond}(A)}{1 - \text{cond}(A)\varepsilon_M} (\varepsilon_M + \varepsilon_M) \approx 2 \text{cond}(A)\varepsilon_M$$

if $\text{cond}(A) \ll 1/\varepsilon_M$. Therefore the unavoidable error is $2 \text{cond}(A)\varepsilon_M$.

Question 2: After we computed \hat{x} how can we check how good our computation was? The obvious thing to check is $\hat{b} := A\hat{x}$ and to compare it with b . The difference $r = \hat{b} - b$ is called the *residual*. As $Ax = b$ and $A\hat{x} = \hat{b}$ we have

$$\frac{\|\hat{x}-x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\hat{b}-b\|}{\|b\|}$$

where $\|\hat{b}-b\|/\|b\|$ is called the *relative residual*. We can compute (or at least estimate) $\text{cond}(A)$, and therefore can obtain an upper bound for the error $\|\hat{x}-x\|/\|x\|$. Actually, we can obtain a slightly better estimate by using

$$\|\hat{x}-x\| = \|A^{-1}(\hat{b}-b)\| \leq \|A^{-1}\| \|\hat{b}-b\| \implies \frac{\|\hat{x}-x\|}{\|\hat{x}\|} \leq \text{cond}(A) \frac{\|\hat{b}-b\|}{\|A\| \|\hat{x}\|}$$

with the *weighted residual* $\rho := \frac{\|\hat{b}-b\|}{\|A\| \|\hat{x}\|}$. Note that $\|\hat{x}-x\|/\|\hat{x}\| \leq \delta$ implies for $\delta < 1$

$$\|\hat{x}-x\| \leq \delta \|x + (\hat{x}-x)\| \leq \delta (\|x\| + \|\hat{x}-x\|) \implies \frac{\|\hat{x}-x\|}{\|x\|} \leq \frac{\delta}{1-\delta}$$

which is the same as δ up to higher order terms $O(\delta^2)$.

If $\|\hat{b}-b\|/\|b\|$ is not much larger than ε_M then the computation was numerically stable: Just perturbing the input slightly from b to \hat{b} and then doing everything else exactly would give the same result \hat{x} .

But it can happen that the relative residual is much larger than ε_M , and yet the computation is numerically stable. We obtain a better way to measure numerical stability by considering perturbations of the matrix A :

Assume we have a computed solution \hat{x} . If we can find a slightly perturbed matrix \tilde{A} such that

$$\frac{\|\tilde{A}-A\|}{\|A\|} \leq \varepsilon, \quad \tilde{A}\hat{x} = b \tag{2}$$

where ε not much larger than ε_M , then the computation is numerically stable: Just perturbing the matrix within the roundoff error and then doing everything exactly gives the same result as our computation.

How can we check whether such a matrix \tilde{A} exists? We again use the “weighted residual”

$$\rho := \frac{\|\hat{b}-b\|}{\|A\| \|\hat{x}\|}$$

Then ρ is the minimal size of a perturbation $\frac{\|\tilde{A}-A\|}{\|A\|}$ giving $\tilde{A}\hat{x} = b$, i.e.,

1. If \hat{x} is the solution of a slightly perturbed problem $\tilde{A}\hat{x} = b$ we have $\frac{\|\tilde{A}-A\|}{\|A\|} \geq \rho$.
2. There is a matrix \tilde{A} with $\frac{\|\tilde{A}-A\|}{\|A\|} = \rho$ such that $\tilde{A}\hat{x} = b$.

Proof:

1. Let $E = \tilde{A} - A$. Then $(A + E)\hat{x} = b$ or $\hat{b} - b = -E\hat{x}$ yielding

$$\|\hat{b} - b\| \leq \|E\| \|\hat{x}\|, \quad \rho = \frac{\|\hat{b} - b\|}{\|A\| \|\hat{x}\|} \leq \frac{\|E\|}{\|A\|}.$$

2. Let $y := b - \hat{b}$. Using Lemma 2 we get a matrix E with $E\hat{x} = y$ and $\|E\| = \frac{\|y\|}{\|\hat{x}\|}$. Then $\tilde{A} := A + E$ satisfies $\tilde{A}\hat{x} = (A + E)\hat{x} = \hat{b} + (b - \hat{b}) = b$ and $\frac{\|E\|}{\|A\|} = \frac{\|b - \hat{b}\|}{\|\hat{x}\| \|A\|} = \rho$.

Summary

- Recommended method for solving linear systems on a computer:
 1. Given A find L, U, p using **Gaussian elimination with pivoting**, choosing the pivot candidate with the largest absolute value.
 2. Solve $Lu = \tilde{b}$ (where $\tilde{b}_i = b_{p_i}$) by forward substitution and $Ux = y$ by back substitution.
- DO NOT compute the inverse matrix A^{-1} . This takes about 3 times as long as computing the LU decomposition.
- The **condition number** $\text{cond}(A) = \|A\| \|A^{-1}\|$ characterizes the sensitivity of the linear system:
 - If $Ax = b$ and $A\hat{x} = \hat{b}$ we have $\frac{\|\hat{x} - x\|}{\|x\|} \leq \text{cond}(A) \frac{\|\hat{b} - b\|}{\|b\|}$.
 - The **unavoidable error** due to the rounding of A and b is approximately $2 \text{cond}(A) \varepsilon_M$.
 - If $\text{cond}(\hat{A}) \geq 1/\varepsilon_M$ then the matrix \hat{A} could be the machine representation of a singular matrix A , and the computed solution is usually meaningless.
 - **To check whether a matrix A is singular, up to a perturbation within machine accuracy:**
 - * DO NOT use the determinant $\det A$
 - * DO NOT use the pivot elements u_{jj} obtained by Gaussian elimination
 - * **DO use the condition number:** $\text{cond}(A) \geq 1/\varepsilon_M \iff$ there exists a singular matrix \tilde{A} with $\frac{\|\tilde{A} - A\|}{\|A\|} \leq \varepsilon_M$
- **Compute an approximation cond1 to the the condition number** $\text{cond}_1(A) = \|A\|_1 \|A^{-1}\|_1$ using **cond1 = norm(A, 1) * invnormest(L, U)** (you need to download `invnormest.m` from course web page). This avoids computing A^{-1} which is very expensive. We have already computed L, U , and `invnormest(L, U)` uses this to solve a few linear systems, yielding a good approximation of $\|A^{-1}\|_1$.
- In order to **check the accuracy of a computed solution** \hat{x} compute the **residual** $r := A\hat{x} - b$ and the **weighted residual** $\rho := \frac{\|r\|}{\|A\| \|\hat{x}\|}$.
 - we get an error bound $\frac{\|\hat{x} - x\|}{\|\hat{x}\|} \leq \text{cond}(A) \rho$
 - **if ρ is not much larger than ε_M** the computation is **numerically stable**: Perturbing A and b with a relative error ε_M would cause the same error.
 - **if ρ is much larger than ε_M** the computation is **numerically unstable**: The error is much larger than the error resulting from the uncertainty of the input values.
We can obtain a more accurate result by **iterative improvement**: Let $r := A\hat{x} - b$ and solve $Ae = r$ using the existing LU decomposition. Then let $\hat{x} := \hat{x} - e$.
- Gaussian elimination with the pivoting strategy of choosing the largest absolute value is in almost all cases numerically stable. We can check this by computing the weighted residual ρ . If ρ is much larger than ε_M we can compute a numerically stable result by using iterative improvement.